

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.




Fig. 1





Fig. 2

Fiber Optic Array Preparation



(B)

BEAD SUBPOPULATION PREPARATION




Fig. 3

(A)




Fig. 4

Fig. 5A






Fig 5B




Polymer Coated Beads in Wells After Air Pulse and Tapping



Before Tapping

Fig. 6A



After Tapping

Fig. 6B





Fig. 7

665070 243360




~~Fig 7~~

~~Fig 7~~ Fig 8

Fig. 9

Porous 3 μ m silica beads high-speed response to
Saturated Toluene




~~scribbles~~

~~scribbles~~

Fig. 10

PMS Beads in Fiber: Response to Methanol (sat)

39 beads, mid-fiber region: centroid ovals, standardized responses



1	2
3	4
5	6
7	8
9	10
11	12
13	14
15	16
17	18
19	20
21	22
23	24
25	26
27	28
29	30
31	32
33	34
35	36
37	38
39	

[Handwritten signature]

[Handwritten mark]



Fig. 11A/11B

PS802 648.c Beads

bead 1

Saturated Toluene

Saturated Methanol




~~663040" E/S/3233~~

~~663040" E/S/3233~~

Fig. 12A/12B

PDPO/psil 9/11



(A)


(B)

mt

mt

Fig. 13

Porous 3 μ m silica beads response to Ethyl Acetate



食言

John R. Smith

[Signature]

Normalized signal-to-noise comparison between bead #1 and summed responses of 39 beads




W. S. S. S. S.



Fig. 15

Signal Enhancement Through Multi-bead Response Summing



Handwritten signature

Handwritten signature

Handwritten signature

"Thick-layer" PS802/Silica 3.2 μ m Beads

- Saturated vapor
- 100X objective
- 50ms exposure

met1	met2	toll	toll2
------	------	------	-------

260.1A

[Signature]

Atkins



Self-Encoding Array with Two Bead Types in Image Guide Wells




Fig. 17

Self-Encoding Array with Two Bead Types in Image Guide Wells

1 2




Fig. 18

Chart4

Self-Encoding Array with Two Bead Types in Image Guide Wells

1 2






Fig.19

Swelling of three different bead types in presence of saturated toluene vapor

PS802 648.c



Poly methyl styrene/
2% divinyl benzene



Poly methyl styrene




Fig. 20

665040" E2523660

Fig 21

Probe	[Cy5]	[Tamra]	[Eu-dye]	Correct Target Identification
HWt	1		0.1	93%
Bglo	0.5		0.05	88%
KWt	0.5		0.005	91%
IL6	0.1		0.1	96%
IL4	0.1		0.005	95%
IFNG		0.4	0.005	95%
IL2		0.04	0.05	98%

Fig 22

1) β -glo (segment of human β -globin)²⁴
TCA ACT TCA TCC ACG TTC ACC

2) IFNG (interferon gamma 1)²⁴
IFNG TGG GTT CTC TTG GCT GTT ACT

3) IL2 (interleukin-2)²⁴
TA CAA GAA TCC CAA ACT CAC CAG

4) IL4 (interleukin-4)²⁴
CC AAC TGC TTC CCC CTC TGT

5) IL6 (interleukin-6)²⁴
GT TGG GTC AGG GGT GGT TAT T

6) K-ras WT²⁷
GGA GCT GGT GGC GTA

7) H-ras WT²⁷
CCG GCG GTG T

8) CFTR (cystic fibrosis exon 11)¹³
CAT TAT ACT TGT AGA G

9) RS53X (cystic fibrosis exon 10)¹³
TGT AGA ATT ATC TTC

10) PAN132¹⁶ (human peripheral lymphocyte)
CCT CTA TAC TTT AAC GTC AAG

11) Schema-2¹⁶
AAG TTT AAC CTA TAC CCT GTC

12) Hakala-1²⁰
CCT ATG ATG AAT ATA G

13) Hakala-2²⁰
AAT ATG ATA ATG GGC T

14) complement to probe 1
TG AAC GTG GAT GAA GTT G

15) complement to probe 2
AG TAA CAG CCA AGA GAA CCC AAA

16) complement to probe 3
CT GGT GAG TTT GGG ATT CTT GTA

17) complement to probe 4
AC AGA GGG GGA AGC AGT TGG

18) complement to probe 5
AA TAA CCA CCC CTG ACC CAA C

19) complement to probe 6
TAC CCC ACC AGC TCC

20) complement to probe 7
ACA CCG CCG G

21) complement to probe 8
CTC TAC AAG TAT AAT G

22) complement to probe 9
GAA GAT GTT AAA GTA TAG AGG

23) complement to probe 10
CTA GAC GTT AAA GTA TAG AGG

24) complement to probe 12
CTA TAT TCA TCA TAG G

25) complement to probe 13
AGG CCA TTA TCA TAT T

3-29-1999 2:00PM

M TUFTS CHEMISTRY 617 627 344

Fig 23

Target Identity	No. of beads in analysis section	[Cy 5]	[Tamra]	[Eu-dye]	Correct Target Identification
2	19		3	0.5	89%
4	15	0.01	0.1	0.1	87%
5	13		0.1	0.1	100%
9	5	0.01			100%
10	14			0.001	86%
11	12		0.1		92%
15	8	0.01	0.1		100%
16	24	0.1			92%
21	21	0.1	3		95%
24	16	0.3	3		94%

665640 " 66564050


Table 4

Sequences	Number of microspheres	Mean background \pm s.d.	Fluorescence after hybridization	Signal
Complementary target	10	997.01 \pm 4.62	1036.94	39.93
	10	1003.46 \pm 6.05	1035.83	32.37
	10	957.44 \pm 5.59	985.25	27.81
	100	977.88 \pm 3.21	1010.74	32.86
Poly A	10	1213.79 \pm 6.33	1221.61	(7.81)
	10	1185.25 \pm 9.39	1194.74	(9.49)
	10	1190.20 \pm 4.85	1198.35	(8.15)
	100	1190.67 \pm 4.05	1199.81	(9.14)
IL2	10	1090.58 \pm 4.97	1096.11	(5.53)
	10	1120.62 \pm 3.09	1113.88	(-6.74)
	10	1101.82 \pm 5.51	1091.28	(-10.55)
	100	1104.36 \pm 1.40	1103.06	(-1.30)

Fig 24

fig 15

CV = 12.4%



CV = Coefficient of variance

$$CV = \frac{\text{St. deviation}}{\text{mean}}$$

Basic idea is that CV decreases
as more bead responses are
summed.

66340" 2.23260


663043 642355

CV = 0.1300
%CV = 13%

%CV = 6.3%

%CV = 4.8%

Fig 25
(Cont.)



CV = 3.5%

2.5%

7.2%